



Spiders

Order: Araneae

The spiders are a large, distinct and widespread group of Arachnids occurring in many types of habitats. Many people think that all spiders are very poisonous; however, although all spiders have venom glands, they very seldom bite people. Most spiders are beneficial because they feed on insects.

Description

A spider's body is divided into two sections: the cephalothorax, which bears the eyes, mouthparts, and legs; and the abdomen, which bears the genital structures, spiracles and anus. Unlike insects, spiders have eight legs and no antennae.

Commonly encountered spiders



Jumping spiders

Small to medium in size, with short legs and stout bodies. The body is hairy and may be brightly colored or iridescent. They are found primarily under stones and in debris.



Crab spiders

As their name suggests, these medium sized spiders are crab-like in appearance, and walk sideways or backwards. These spiders spin no webs, but forage for their prey or wait in ambush for it. One species can change color depending upon the color of the flower it is resting on.




Nursery Web spiders and Fishing spiders

These may be quite large, some having a leg spread of three inches. Many of them live near water, walking over the surface and diving beneath it. They feed on aquatic insects and even small fish. These spiders are rovers, and the female spins a web only for the young. The female carries the egg sac underneath her until the young spiders are ready to hatch, at which time she ties it to a plant and wraps leaves around it.




Orb Weavers

Included in this group is the common garden spider. Many are brightly colored, being black and yellow, or black and red. These spiders are usually found resting head downward near the center of their large orb web.



Ground spiders and Wolf spiders

Most of these common spiders are quite large, and dark brown in color. They may be found running along the ground chasing their prey. The females carry the egg sac, and when the young hatch, they are carried on the female's back for a few days.



Harvestman or Daddy-Longlegs

Although these animals are not true spiders, but rather a separate order of Arachnids, the Harvestmen and Daddy-Longlegs deserve mention here because they are found in our gardens.

These Arachnids have a small oval body, and 8 extremely long legs (only 4 are shown in this drawing). They feed on plant juices and dead insects.

Some spiders considered to be dangerous



Photo by Ronald F. Billings,
Texas Forest Service, from Bugwood.org

Tarantulas

These spiders are feared because of their size and formidable appearance. The bodies may get up to 3 to 3½ inches in length. Actually, they are sluggish spiders, and will attack only when they are teased to an extreme. Many are kept for pets. The larger species can inflict a painful wound with their powerful jaws. Tarantulas are desert spiders, and are not found in Northeastern States.



Photo by Sturgis McKeever,
Georgia Southern University, from Bugwood.org

Black Widow spider

(*Latrodectus* spp.; *L. mactans* is now thought to be 3 species)

The black widow is timid by nature; however, when accidentally squeezed, it will bite in self-defense. The venom is highly toxic. The bite causes extreme pain, which usually extends to the abdominal muscles. The danger exists because the black widow seeks out dark places in which to construct a retreat, often living in close proximity to people. Female black widow spiders are about a half inch long, with 1 1/2 inch leg spread; males are half that size.



Photo by Lisa Ames,
University of Georgia, from Bugwood.org

Brown Recluse spider

(*Loxosceles reclusa*)

This spider has a dark violin-shaped marking on its back. Its bite can cause sores that are slow to heal. These range from western Georgia northward into Kentucky and westward to Kansas and Texas. They are not found in the Northeast, but since these spiders are easily transported, it is wise to check vacation gear when returning from a southern and western vacation.

The spiders hide in dark corners and are generally shy. They vary in size, with a body length around 3/8 inch, and leg spread about an inch. Recluse spiders have 6 eyes (not 8 like many other spiders).



Photo by Joseph Berger,
from Bugwood.org

Yellow Sac spider

(*Cheiracanthium* spp.; Family Clubionidae)

Several species of sac spiders can be found indoors. They are small, with a body length generally 3/8 inch, and leg span about 1 inch. These spiders can cause painful bites. In the past they were also implicated in causing necrotic lesions, but recent information indicates that they do not cause these.

(see: <http://www.ajtmh.org/cgi/content/abstract/74/6/1043>)

Further information on venomous spiders and spider bite symptoms is available from the Centers for Disease Control and Prevention: <http://www.cdc.gov/niosh/topics/spiders/>

Management

Although most spiders are beneficial, they can become a nuisance when found in the home. Keep spiders out with proper screening and weather stripping. Remove indoor webs with a broom or dustmop, and discard outdoors. Vacuum up or remove spiders with a dustcloth. It may take some time, but eventually spiders will be controlled. Additionally, control indoor insects on which spiders feed.

If necessary, a pesticide labelled for indoor use on spiders may be used for control. Spray in corners of rooms, at both floor and ceiling levels.

Outdoors, keeping vegetation away from the side of the house can help to discourage spiders from constructing webs. Using yellow light bulbs at night if lights are left on outdoors can also help – they attract fewer insects, so there is less food available for spiders, which will move elsewhere.

Prepared 1995 by Carolyn Klass, Senior Extension Associate, Dept. of Entomology, Cornell University
Updated 2012

This publication contains pesticide recommendations. Changes in pesticide regulations occur constantly and human errors are still possible. Some materials mentioned may no longer be available and some uses may no longer be legal. All pesticides distributed, sold or applied in New York State must be registered with the New York State Department of Environmental Conservation (DEC). Questions concerning the legality and/or registration status for pesticide use in New York State should be directed to the appropriate Cornell Cooperative Extension Specialist or your regional DEC office. READ THE LABEL BEFORE APPLYING ANY PESTICIDE.

<http://idl.entomology.cornell.edu>